

# Calculus for mathematicians

D. J. Bernstein, University of Illinois at Chicago, [djb@pobox.com](mailto:djb@pobox.com)

19970403, draft 5

## 1. Introduction

This booklet presents the main concepts, theorems, and techniques of single-variable calculus. It differs from a typical undergraduate real analysis text in that (1) it focuses purely on calculus, not on developing topology and analysis for their own sake; (2) it's short.

**Notation and terminology.** The reader must be comfortable with *functions*, not just numbers, as objects of study. I use the notation  $x \mapsto x^2$  for the function that takes  $x$  to  $x^2$ ; thus  $(x \mapsto x^2)(3) = 9$ . In general  $f = (t \mapsto f(t))$  for any function  $f$ .

An **open ball around  $c$**  means an interval  $\text{Ball}(c, h) = \{x : |x - c| < h\}$  for some positive real number  $h$ . The intersection of two open balls around  $c$  is another open ball around  $c$ .

If  $S$  is a set, and  $f(x)$  is defined for all  $x \in S$ , then  $f(S)$  is defined as  $\{f(x) : x \in S\}$ .

## Part I. Continuity

### 2. Continuous functions

**Definition 2.1.** *Let  $f$  be a function defined at  $c$ . Then  $f$  is **continuous at  $c$**  if, for any open ball  $F$  around  $f(c)$ , there is an open ball  $B$  around  $c$  such that  $f(B) \subseteq F$ .*

In other words, if  $f$  is continuous at  $c$ , and  $F$  is an open ball around  $f(c)$ , then there is some  $h > 0$  such that  $f(x) \in F$  for all  $x$  with  $|x - c| < h$ .

Example: The function  $x \mapsto 3x$  is continuous—i.e., continuous at  $c$  for every  $c$ . Indeed,  $\text{Ball}(3c, \epsilon)$  contains  $(x \mapsto 3x)(\text{Ball}(c, \epsilon/3))$ , because  $|x - c| < \epsilon/3$  implies  $|3x - 3c| < \epsilon$ .

Another example: If  $f(x) = 3$  for  $x < 2$  and  $f(x) = 5$  for  $x \geq 2$ , then  $f$  is not continuous at 2. Indeed, consider the open ball  $F = \text{Ball}(5, 1)$ . If  $B$  is any open ball around 2, then  $B$  contains numbers smaller than 2, so  $3 \in f(B)$ ; thus  $f(B)$  is not contained in  $F$ .

**Theorem 2.2.** *Let  $f$  and  $g$  be functions continuous at  $c$ . Assume that  $f(x) = g(x)$  for all  $x \neq c$  such that  $f(x)$  and  $g(x)$  are both defined. Then  $f(c) = g(c)$ .*

**Proof.** I will show that  $|f(c) - g(c)| < 2\epsilon$  for any  $\epsilon > 0$ . Write  $F = \text{Ball}(f(c), \epsilon)$  and  $G = \text{Ball}(g(c), \epsilon)$ . By continuity of  $f$  and  $g$ , there are balls  $A$  and  $B$  around  $c$  such that  $f(A) \subseteq F$  and  $g(B) \subseteq G$ . Find a point  $x \neq c$  contained in both  $A$  and  $B$ . By construction  $f(x) \in F$  and  $f(x) = g(x) \in G$ , so  $|f(c) - g(c)| \leq |f(x) - f(c)| + |f(x) - g(c)| < 2\epsilon$  as claimed.  $\square$

### 3. Continuity of sums, products, and compositions

**Theorem 3.1.** *Let  $f$  and  $g$  be functions continuous at  $c$ . Define  $h = f + g$ . Then  $h$  is continuous at  $c$ .*

**Proof.** Given a ball  $H = \text{Ball}(h(c), \epsilon)$ , consider the balls  $F = \text{Ball}(f(c), \epsilon/2)$  and  $G = \text{Ball}(g(c), \epsilon/2)$ . By continuity of  $f$  and  $g$ , there are open balls  $A$  and  $B$  around  $c$  such that  $f(A) \subseteq F$  and  $g(B) \subseteq G$ . Define  $D = A \cap B$ ;  $D$  is an open ball around  $c$ . If  $x \in D$  then  $f(x) \in F$  and  $g(x) \in G$  so  $h(x) = f(x) + g(x) \in H$ . Thus  $h(D) \subseteq H$ .  $\square$

**Theorem 3.2.** *Let  $f$  and  $g$  be functions continuous at  $c$ . Define  $h = fg$ . Then  $h$  is continuous at  $c$ .*

**Proof.** Define  $L = f(c)$  and  $M = g(c)$ , so that  $LM = h(c)$ . Given an open ball  $H = \text{Ball}(LM, \epsilon)$ , I will find an open ball  $D$  around  $c$  so that  $h(D) \subseteq H$ .

If  $L = M = 0$ , take the intersection of open balls where  $|f(x)| < \epsilon$  and  $|g(x)| < 1$ . Then  $|h(x)| < \epsilon$ .

If  $L = 0$  and  $M \neq 0$ , take the intersection of open balls where  $|f(x)| < \epsilon/(2|M|)$  and  $|g(x) - M| < |M|$ . Then  $|g(x)| < 2|M|$  so  $|h(x)| < \epsilon$ . Similarly if  $L \neq 0$  and  $M = 0$ .

If  $L \neq 0$  and  $M \neq 0$ , take the intersection of open balls where  $|f(x) - L| < \epsilon/(4|M|)$ ,  $|g(x) - M| < \epsilon/(2|L|)$ , and  $|g(x) - M| < |M|$ . Then  $|g(x)| < 2|M|$  so  $|h(x) - LM| = |g(x)(f(x) - L) + L(g(x) - M)| < 2|M|(\epsilon/(4|M|)) + |L|(\epsilon/(2|L|)) = \epsilon$ .  $\square$

**Theorem 3.3.** *Let  $g$  be a function continuous at  $c$ . Let  $f$  be a function continuous at  $g(c)$ . Define  $h = (x \mapsto f(g(x)))$ . Then  $h$  is continuous at  $c$ .*

For example,  $x \mapsto \cos 2x$  is continuous, since  $x \mapsto 2x$  and  $y \mapsto \cos y$  are continuous,

**Proof.** Let  $F$  be an open ball around  $h(c) = f(g(c))$ . By continuity of  $f$ , there is some open ball  $G$  around  $g(c)$  with  $f(G) \subseteq F$ . By continuity of  $g$ , there is some open ball  $B$  around  $c$  with  $g(B) \subseteq G$ . Finally  $h(B) = f(g(B)) \subseteq f(G) \subseteq F$ .  $\square$

### 4. Continuity of simple functions

**Theorem 4.1.**  *$x \mapsto b$  is continuous at  $c$ , for any  $b$  and  $c$ .*

**Proof.**  $\text{Ball}(b, h)$  contains  $(x \mapsto b)(D)$  for any open ball  $D$ .  $\square$

**Theorem 4.2.**  *$x \mapsto x$  is continuous at  $c$ , for any  $c$ .*

**Proof.**  $\text{Ball}(c, h)$  contains  $(x \mapsto x)(\text{Ball}(c, h))$ .  $\square$

By Theorems 3.2 and 4.2,  $x \mapsto x^2$  is continuous;  $x \mapsto x^3$  is continuous; in general  $x \mapsto x^n$  is continuous for any positive integer  $n$ . Thus, by Theorems 3.1, 3.2, and 4.1, any polynomial function  $x \mapsto c_0 + c_1x + \cdots + c_nx^n$  is continuous.

The function  $x \mapsto 1/x$  is continuous at  $c$  for  $c \neq 0$ . (It's not even defined at 0, so it can't be continuous there.) By Theorem 3.3,  $x \mapsto 1/f(x)$  is continuous whenever  $f$  is continuous and nonzero. For example,  $x \mapsto x^n$  is continuous except at 0 when  $n$  is a negative integer.

## Part II. Derivatives

### 5. Differentiable functions

**Definition 5.1.** Let  $f$  be a function defined at  $c$ . Then  $f$  is **differentiable at  $c$**  if there is a function  $f_1$ , continuous at  $c$ , such that  $f = (x \mapsto f(c) + (x - c)f_1(x))$ .

**Definition 5.2.** Let  $f$  be a function defined at  $c$ . Then  $f$  has **derivative  $d$  at  $c$**  if there is a function  $f_1$ , continuous at  $c$ , such that  $f = (x \mapsto f(c) + (x - c)f_1(x))$  and  $f_1(c) = d$ .

By Theorem 2.2, there is at most one continuous function  $f_1$  satisfying  $f_1(x) = (f(x) - f(c))/(x - c)$  for all  $x \neq c$ , so  $f$  has at most one derivative at  $c$ , called **the derivative of  $f$  at  $c$** . The derivative of  $f$  at  $c$  is written  $f'(c)$ . The **derivative of  $f$** , written  $f'$ , is the function  $c \mapsto f'(c)$ .

For example, consider the function  $f = (x \mapsto x^2)$ . Here  $f(x) = f(3) + (x - 3)f_1(x)$  with  $f_1 = (x \mapsto x + 3)$ . The function  $f_1$  is continuous at 3, so  $f$  is differentiable at 3; its derivative at 3 is  $f_1(3) = 6$ . In general  $f'(c) = 2c$ .

**Theorem 5.3.** Let  $f$  be a function. If  $f$  is differentiable at  $c$  then  $f$  is continuous at  $c$ .

**Proof.** By definition of differentiability, there is a function  $f_1$ , continuous at  $c$ , with  $f = (x \mapsto f(c) + (x - c)f_1(x))$ . Apply Theorems 3.1, 3.2, 4.1, and 4.2.  $\square$

### 6. Derivatives of sums, products, and compositions

**Theorem 6.1.** Let  $f$  and  $g$  be functions. Define  $h = f + g$ . If  $f$  and  $g$  are differentiable at  $c$  then  $h$  is differentiable at  $c$ . Furthermore  $h'(c) = f'(c) + g'(c)$ .

In short  $(f + g)' = f' + g'$  if the right side is defined. This is the **sum rule**.

**Proof.** Say  $f(x) = f(c) + (x - c)f_1(x)$  and  $g(x) = g(c) + (x - c)g_1(x)$  with  $f_1$  and  $g_1$  continuous at  $c$ . Define  $h_1 = f_1 + g_1$ ; then  $h_1$  is continuous at  $c$  by Theorem 3.1, and  $h(x) = h(c) + (x - c)h_1(x)$ , so  $h$  is differentiable at  $c$ . Finally  $h'(c) = h_1(c) = f_1(c) + g_1(c) = f'(c) + g'(c)$ .  $\square$

**Theorem 6.2.** Let  $f$  and  $g$  be functions. Define  $h = fg$ . If  $f$  and  $g$  are differentiable at  $c$  then  $h$  is differentiable at  $c$ . Furthermore  $h'(c) = f'(c)g(c) + f(c)g'(c)$ .

In short  $(fg)' = f'g + fg'$  if the right side is defined. This is the **product rule**.

**Proof.** Say  $f(x) = f(c) + (x - c)f_1(x)$  and  $g(x) = g(c) + (x - c)g_1(x)$  with  $f_1$  and  $g_1$  continuous at  $c$ . Then  $h(x) = h(c) + (x - c)h_1(x)$  where  $h_1(x) = f_1(x)g(x) + f(c)g_1(x)$ . This function  $h_1$  is continuous at  $c$  by Theorems 3.1, 3.2, 4.1, and 5.3, so  $h$  is differentiable at  $c$ , with derivative  $h_1(c) = f_1(c)g(c) + f(c)g_1(c) = f'(c)g(c) + f(c)g'(c)$ .  $\square$

**Theorem 6.3.** *Let  $f$  and  $g$  be functions. Define  $h = (x \mapsto f(g(x)))$ . If  $g$  is differentiable at  $c$ , and  $f$  is differentiable at  $g(c)$ , then  $h$  is differentiable at  $c$ . Furthermore  $h'(c) = f'(g(c))g'(c)$ .*

In short  $(f \circ g)' = (f' \circ g)g'$  if the right side is defined. This is the **chain rule**.

**Proof.** Write  $b = g(c)$ . Say  $f(x) = f(b) + (x - b)f_1(x)$  and  $g(x) = b + (x - c)g_1(x)$  with  $f_1$  continuous at  $b$  and  $g_1$  continuous at  $c$ . Now  $h(x) = f(g(x)) = f(b) + (g(x) - b)f_1(g(x)) = f(b) + (x - c)g_1(x)f_1(g(x))$ . Thus  $h(x) = h(c) + (x - c)h_1(x)$  where  $h_1(x) = g_1(x)f_1(g(x))$ . Finally  $h_1$  is continuous at  $c$  by Theorems 3.3, 3.2, and 5.3, so  $h$  is differentiable at  $c$ , with derivative  $h_1(c) = g_1(c)f_1(g(c)) = g'(c)f'(g(c))$ .  $\square$

## 7. Derivatives of simple functions

A constant function, such as  $x \mapsto 17$ , has derivative  $c \mapsto 0$ , since  $17 = 17 + (x - c)0$ .

The identity function  $x \mapsto x$  has derivative  $c \mapsto 1$ , since  $x = c + (x - c)1$ .

In general, for any positive integer  $n$ , the function  $x \mapsto x^n$  has derivative  $c \mapsto nc^{n-1}$ , since  $x^n = c^n + (x - c)(x^{n-1} + cx^{n-2} + \dots + c^{n-1})$ .

The function  $x \mapsto 1/x$ , defined for nonzero inputs, has derivative  $c \mapsto -1/c^2$ . Indeed,  $1/x = 1/c + (x - c)(-1/cx)$ , and  $x \mapsto -1/cx$  is continuous at  $c$  with value  $-1/c^2$ .

Now the chain rule, with  $f = (x \mapsto 1/x)$ , states that  $1/g$  has derivative  $-g'/g^2$  at any point  $c$  where  $g(c) \neq 0$ . In particular, for any negative integer  $n$ ,  $x \mapsto x^n$  has derivative  $c \mapsto nc^{n-1}$ .

Finally, the product rule implies that  $h/g$  has derivative  $(gh' - hg')/g^2$  at any point  $c$  where  $g(c) \neq 0$ ; this is the **quotient rule**.

## Part III. Completeness and its consequences

### 8. Completeness of the real numbers

**Definition 8.1.** *Let  $S$  be a set of real numbers. A real number  $c$  is an **upper bound** for  $S$  if  $x \leq c$  for all  $x \in S$ .*

For example, any number  $c \geq \pi$  is an upper bound for the set  $\{3, 3.1, 3.14, 3.141, \dots\}$ . The smallest upper bound is  $\pi$ .

The real numbers are **complete**: if  $S$  is a nonempty set, and there is an upper bound for  $S$ , then there is a smallest upper bound for  $S$ . The smallest upper bound is unique; it is called the **supremum** of  $S$ , written  $\sup S$ .

## 9. The intermediate-value theorem

**Theorem 9.1.** *Let  $f$  be a continuous real-valued function. Let  $y$  be a real number. Let  $b \leq c$  be real numbers with  $f(b) \leq y \leq f(c)$ . Then  $f(x) = y$  for some  $x \in [b, c]$ .*

Here  $[b, c]$  means  $\{x : b \leq x \leq c\}$ . For example, if  $f(3) = -5$  and  $f(4) = 7$ , and  $f$  is continuous, then  $f$  must have a root between 3 and 4.

**Proof.** Define  $S = \{x \in [b, c] : f(x) \leq y\}$ .  $S$  is nonempty, because it contains  $b$ , and it has an upper bound, namely  $c$ , so it has a smallest upper bound, say  $u$ .

Suppose  $f(u) > y$ . By continuity, there is an open ball  $D$  around  $u$  such that  $f(x) > y$  for  $x \in D$ . Pick any  $t \in D$  with  $t < u$ . If  $x \in [t, u]$  then  $x \in D$  so  $f(x) > y$  so  $x \notin S$ . Thus  $t$  is an upper bound for  $S$ —but  $u$  is the smallest upper bound. Contradiction.

Suppose  $f(u) < y$ . Then  $u \neq c$  so  $u < c$ . By continuity, there is an open ball  $D$  around  $u$  such that  $f(x) < y$  for  $x \in D$ . Pick any  $x \in D$  with  $u < x < c$ ; then  $f(x) < y$ . But  $x \notin S$  since  $u$  is an upper bound for  $S$ ; so  $f(x) > y$ . Contradiction.  $\square$

## 10. The maximum-value theorem

**Theorem 10.1.** *Let  $f$  be a continuous real-valued function. Let  $b \leq c$  be real numbers. Then there is an upper bound for  $f([b, c])$ .*

**Proof.** Let  $S$  be the set of  $x \in [b, c]$  such that  $f([b, x])$  is bounded—i.e., has an upper bound.  $S$  is nonempty, because it contains  $b$ . Define  $u = \sup S$ .

By continuity, there is an open ball  $D$  around  $u$  such that  $f(D) \subseteq \text{Ball}(f(u), 1)$ . Select  $t \in D$  with  $t < u$ ; then  $t$  is not an upper bound for  $S$ , so there is some  $x \in S$  with  $t < x \leq u$ . Now  $f([b, x])$  and  $f([x, u]) \subseteq f(D)$  are bounded, so  $f([b, u])$  is bounded.

Suppose  $u < c$ . Select  $v \in D$  with  $u < v < c$ . Then  $f([u, v])$  is bounded, so  $v \in S$ . Contradiction. Hence  $u = c$ , and  $f([b, c]) = f([b, u])$  is bounded.  $\square$

**Theorem 10.2.** *Let  $f$  be a continuous real-valued function. Let  $b \leq c$  be real numbers. Then there is some  $u \in [b, c]$  such that, for all  $z \in [b, c]$ ,  $f(u) \geq f(z)$ .*

This is the **maximum-value theorem**: a continuous function on a closed interval achieves a maximum. The same is not true for open intervals: consider  $1/x$  for  $0 < x < 1$ .

**Proof.** By Theorem 10.1, there is an upper bound for  $f([b, c])$ . Define  $M = \sup f([b, c])$ .

Let  $S$  be the set of  $x \in [b, c]$  such that  $\sup f([x, c]) = M$ . Then  $b \in S$ . Define  $u = \sup S$ .

Suppose  $f(u) < M$ . By continuity there is an open ball  $D$  around  $u$  such that  $f(D) \subseteq \text{Ball}(f(u), (M - f(u))/2)$ ; then  $\sup f(D) < M$ . Select  $t \in D$  with  $t < u$ ; then  $t$  is not an upper bound for  $S$ , so there is some  $x \in S$  with  $t < x \leq u$ . Then  $\sup f([x, c]) = M$ , but  $\sup f([x, u]) < M$ , so  $u < c$ . Select  $v \in D$  with  $u < v < c$ . Then  $\sup f([x, v]) < M$ , so  $\sup f([v, c]) = M$ , so  $v \in S$ . Contradiction. Hence  $f(u) = M = \sup f([b, c])$ .  $\square$

**Theorem 10.3.** *Let  $f$  be a continuous real-valued function. Let  $b \leq c$  be real numbers. Then there is some  $u \in [b, c]$  such that, for all  $x \in [b, c]$ ,  $f(u) \leq f(x)$ .*

**Proof.** Apply Theorem 10.2 to  $-f$ . □

## Part IV. The mean-value theorem

### 11. Fermat's principle

**Theorem 11.1.** *Let  $f$  be a real-valued function differentiable at  $t$ . Assume that  $f(t) \geq f(x)$  for all  $x$  in an open ball  $B$  around  $t$ . Then  $f'(t) = 0$ .*

**Proof.** By assumption  $f(x) = f(t) + (x - t)f_1(x)$  where  $f_1$  is continuous at  $t$ . Suppose  $f_1(t) > 0$ . Then  $f_1(x) > 0$  for all  $x$  in an open ball  $D$  around  $t$ . Pick  $x > t$  in both  $B$  and  $D$ ; then  $f(t) \geq f(x) = f(t) + (x - t)f_1(x) > f(t)$ . Contradiction. Thus  $f_1(t) \leq 0$ . Similarly  $f_1(t) \geq 0$ . Hence  $f'(t) = f_1(t) = 0$ . □

**Theorem 11.2.** *Let  $f$  be a real-valued function differentiable at  $t$ . Assume that  $f(t) \leq f(x)$  for all  $x$  in an open ball  $B$  around  $t$ . Then  $f'(t) = 0$ .*

**Proof.** Apply Theorem 11.1 to  $-f$ . □

### 12. Rolle's theorem

**Theorem 12.1.** *Let  $f$  be a differentiable real-valued function. Let  $b < c$  be real numbers. If  $f(b) = f(c)$  then there is some  $x$  with  $b < x < c$  such that  $f'(x) = 0$ .*

**Proof.** By Theorem 10.2, there is some  $t \in [b, c]$  such that  $f$ 's maximum value on  $[b, c]$  is achieved at  $t$ . If  $f(t) > f(b)$  then  $t \neq b$  and  $t \neq c$ , so there is an open ball  $B$  around  $t$  such that  $B \subseteq [b, c]$ . By Theorem 11.1,  $f'(t) = 0$ .

Similarly, by Theorem 10.3, there is some  $u \in [b, c]$  such that  $f$  achieves its minimum at  $u$ . If  $f(u) < f(b)$  then  $f'(u) = 0$  as above.

The only remaining case is that  $f(t) \leq f(b)$  and  $f(u) \geq f(b)$ . Then  $f(b)$  is both the maximum and the minimum value of  $f$  on  $[b, c]$ ; i.e.,  $f$  is constant on  $[b, c]$ . Hence  $f'(x) = 0$  for any  $x$  between  $b$  and  $c$ . □

### 13. The mean-value theorem

**Theorem 13.1.** *Let  $f$  be a differentiable real-valued function. Let  $b < c$  be real numbers. Then there is some  $x$  with  $b < x < c$  such that  $f(c) - f(b) = f'(x)(c - b)$ .*

This is the **mean-value theorem**. The terminology “mean value” comes from the fundamental theorem of calculus, which can be interpreted as saying that  $(f(c) - f(b))/(c - b)$  is the average (“mean”) value of  $f'(x)$  for  $x \in [b, c]$ . See Theorem 16.1.

**Proof.** Define  $g(x) = (c - b)f(x) - (x - b)(f(c) - f(b))$ . Then  $g$  is differentiable, and  $g(b) = (c - b)f(b) = (c - b)f(c) - (c - b)(f(c) - f(b)) = g(c)$ . By Theorem 12.1,  $g'(x) = 0$  for some  $x$  between  $b$  and  $c$ . But  $g'(x) = (c - b)f'(x) - (f(c) - f(b))$ .  $\square$

**Theorem 13.2.** *Let  $f$  be a differentiable real-valued function. If  $f'(x) = 0$  for all  $x$  then  $f$  is constant.*

More generally, two functions with the same derivative must differ by a constant.

**Proof.** Pick any real numbers  $b < c$ . By Theorem 13.1, there is some  $x$  such that  $f(c) - f(b) = f'(x)(c - b) = 0$ , so  $f(c) = f(b)$ .  $\square$

## Part V. Integration

### 14. Tagged divisions and gauges

**Definition 14.1.** *Let  $b \leq c$  be real numbers. Let  $x_0, x_1, \dots, x_n$  and  $t_1, \dots, t_n$  be real numbers. Then  $x_0, t_1, x_1, \dots, t_n, x_n$  is a **tagged division of  $[b, c]$**  if  $b = x_0 \leq t_1 \leq x_1 \leq t_2 \leq \dots \leq x_{n-1} \leq t_n \leq x_n = c$ .*

The idea is that  $[b, c]$  is divided into the intervals  $[x_0, x_1], [x_1, x_2], \dots, [x_{n-1}, x_n]$ ; in each interval  $[x_{k-1}, x_k]$  there is a *tag*  $t_k$ . For example, consider the tagged division  $0, 1, 4, 5, 6, 6, 7$  of  $[0, 7]$ ; here the intervals are  $[0, 4], [4, 6], [6, 7]$ , with tags  $1, 5, 6$  respectively.

**Definition 14.2.** *Let  $b \leq c$  be real numbers. A **gauge on  $[b, c]$**  is a function assigning to each point  $t \in [b, c]$  an open interval containing  $t$ .*

For example, given  $\epsilon > 0$ , the function  $t \mapsto \text{Ball}(t, \epsilon)$  is a gauge on any interval.

**Definition 14.3.** *Let  $b \leq c$  be real numbers. Let  $G$  be a gauge on  $[b, c]$ . A tagged division  $x_0, t_1, x_1, \dots, t_n, x_n$  of  $[b, c]$  is **inside  $G$**  if  $[x_{k-1}, x_k] \subset G(t_k)$  for every  $k$ .*

**Theorem 14.4.** *Let  $b \leq c$  be real numbers. Let  $G$  be a gauge on  $[b, c]$ . Then there is a tagged division of  $[b, c]$  inside  $G$ .*

**Proof.** Let  $S$  be the set of  $x \in [b, c]$  such that there is a tagged division of  $[b, x]$  inside  $G$ .  $S$  is nonempty:  $b, b, b$  is a tagged division of  $[b, b]$  inside  $G$ , so  $b \in S$ . Also,  $c$  is an upper bound for  $S$ . Thus there is a smallest upper bound for  $S$ , say  $y$ .

Select  $v \in G(y)$  such that  $v < y$ . Then  $v$  is not an upper bound for  $S$ , so there is some  $x > v$  with  $x \in S$ . Let  $x_0, t_1, x_1, \dots, t_n, x_n$  be a tagged division of  $[b, x]$  inside  $G$ .

Suppose  $y < c$ . Pick  $z \in G(y)$  with  $y < z \leq c$ . Then  $[x_n, z] = [x, z] \subset [v, z] \subset G(y)$ , so  $x_0, t_1, x_1, \dots, t_n, x_n, y, z$  is a tagged division of  $[b, z]$  inside  $G$ . Thus  $z \in S$ ; but  $z > y$ , and  $y$  is an upper bound for  $S$ . Contradiction.

Thus  $y = c$ . Finally  $x_0, t_1, x_1, \dots, t_n, x_n, y, y$  is a tagged division of  $[b, c]$  inside  $G$ .  $\square$

## 15. The definite integral

**Definition 15.1.** Let  $b \leq c$  be real numbers. Let  $x_0, t_1, x_1, \dots, t_n, x_n$  be a tagged division of  $[b, c]$ . Let  $f$  be a function defined on  $[b, c]$ . The **Riemann sum for  $f$  on  $x_0, t_1, x_1, \dots, t_n, x_n$**  is  $(x_1 - x_0)f(t_1) + \dots + (x_n - x_{n-1})f(t_n)$ .

For example, the Riemann sum for  $f$  on  $0, 1, 4, 5, 6, 6, 7$  is  $(4 - 0)f(1) + (6 - 4)f(5) + (7 - 6)f(6)$ . This may be visualized as the sum of areas of three rectangles: one stretching from 0 to 4 horizontally with height  $f(1)$ , another from 4 to 6 with height  $f(5)$ , and another from 6 to 7 with height  $f(6)$ .

**Definition 15.2.** Let  $b \leq c$  be real numbers. Let  $f$  be a function defined on  $[b, c]$ . Let  $I$  be a number. Then  $f$  **has integral  $I$  on  $[b, c]$**  if, for every open ball  $E$  around  $I$ , there is a gauge  $G$  on  $[b, c]$  such that  $E$  contains the Riemann sum for  $f$  on any tagged division of  $[b, c]$  inside  $G$ .

**Theorem 15.3.** Let  $b \leq c$  be real numbers. Let  $f$  be a function. If  $f$  has integral  $I$  on  $[b, c]$  and  $f$  has integral  $J$  on  $[b, c]$  then  $I = J$ .

Thus there is at most one number  $I$  such that  $f$  has integral  $I$  on  $[b, c]$ . If this number exists, it is called the **integral of  $f$  from  $b$  to  $c$** , written  $\int_b^c f$ .

**Proof.** I will show that  $|I - J| < 2\epsilon$  for any  $\epsilon > 0$ .

By definition of integral, there is a gauge  $G$  on  $[b, c]$  such that  $\text{Ball}(I, \epsilon)$  contains the Riemann sum for  $f$  on any tagged division of  $[b, c]$  inside  $G$ .

Similarly, there is a gauge  $H$  on  $[b, c]$  such that  $\text{Ball}(J, \epsilon)$  contains the Riemann sum for  $f$  on any tagged division of  $[b, c]$  inside  $H$ .

Define  $F(t)$  as the intersection of  $G(t)$  and  $H(t)$ . Then  $F$  is a gauge on  $[b, c]$ . By Theorem 14.4, there is a tagged division  $x_0, \dots, x_n$  of  $[b, c]$  inside  $F$ .

Let  $R$  be the Riemann sum for  $f$  on  $x_0, \dots, x_n$ . Observe that  $x_0, \dots, x_n$  is inside both  $G$  and  $H$ , so  $R \in \text{Ball}(I, \epsilon)$  and  $R \in \text{Ball}(J, \epsilon)$ . Hence  $|I - J| \leq |I - R| + |R - J| < 2\epsilon$ .  $\square$

## 16. The fundamental theorem of calculus

**Theorem 16.1.** Let  $f$  be a differentiable function. Let  $b \leq c$  be real numbers. Then  $f(c) - f(b) = \int_b^c f'$ .



**Proof.** Pick  $\epsilon > 0$ . I will construct a gauge  $G$  such that  $\text{Ball}(f(c) - f(b), \epsilon(c - b + 1))$  contains the Riemann sum for  $f'$  on any tagged division of  $[b, c]$  inside  $G$ .

Fix  $t \in [b, c]$ . Since  $f$  is differentiable at  $t$ , there is a function  $f_1$ , continuous at  $t$ , such that  $f(x) = f(t) + (x - t)f_1(x)$ . By definition of continuity,  $f_1(x)$  is within  $\epsilon$  of  $f_1(t) = f'(t)$  for all  $x$  in some open ball around  $t$ . Define  $G(t)$  as the union of all such balls. Then  $G$  is a gauge on  $[b, c]$ .

Observe that if  $x, y \in G(t)$ , with  $x \leq t \leq y$ , then  $(y - x)f'(t)$  is within  $\epsilon(y - x)$  of  $f(y) - f(x)$ . Indeed,  $|f_1(x) - f'(t)| < \epsilon$  by definition of  $G$ , and  $f(x) - f(t) = (x - t)f_1(x)$ , so

$$|f(x) - f(t) - (x - t)f'(t)| = |(x - t)(f_1(x) - f'(t))| \leq \epsilon|x - t|.$$

Similarly  $|f(y) - f(t) - (y - t)f'(t)| \leq \epsilon|y - t|$ . Thus  $|f(y) - f(x) - (y - x)f'(t)| \leq \epsilon(|y - t| + |x - t|)$ ; and  $|y - t| + |x - t| = y - x$ .

Finally, say  $x_0, t_1, x_1, \dots, t_n, x_n$  is a tagged division of  $[b, c]$  inside  $G$ . Then  $x_{k-1}, x_k \in G(t_k)$ , with  $x_{k-1} \leq t_k \leq x_k$ , so  $(x_k - x_{k-1})f'(t_k)$  is within  $\epsilon(x_k - x_{k-1})$  of  $f(x_k) - f(x_{k-1})$  as above. Thus the Riemann sum for  $f'$  on  $x_0, t_1, x_1, \dots, t_n, x_n$  is within

$$\sum_{1 \leq k \leq n} \epsilon(x_k - x_{k-1}) = \epsilon(x_n - x_0) = \epsilon(c - b) < \epsilon(c - b + 1)$$

of

$$\sum_{1 \leq k \leq n} (f(x_k) - f(x_{k-1})) = f(x_n) - f(x_0) = f(c) - f(b)$$

as claimed. □

## 17. Integration rules

**Theorem 17.1.** *Let  $f$  be a function. Let  $b \leq c$  be real numbers. If  $\int_b^c f = I$  then  $af$  has integral  $aI$  on  $[b, c]$  for any real number  $a$ .*

In short  $\int_b^c af = a \int_b^c f$  if the right side is defined.

**Proof.** Pick  $\epsilon > 0$ . Since  $\int_b^c f = I$ , there is a gauge  $G$  on  $[b, c]$  such that  $\text{Ball}(I, \epsilon)$  contains the Riemann sum for  $f$  on any tagged division of  $[b, c]$  inside  $G$ . The Riemann sum for  $af$  is exactly  $a$  times the Riemann sum for  $f$ , so it is inside  $\text{Ball}(aI, |a|\epsilon)$  for  $a \neq 0$  or  $\text{Ball}(0, \epsilon)$  for  $a = 0$ . □

**Theorem 17.2.** *Let  $f$  and  $g$  be functions. Let  $b \leq c$  be real numbers. If  $\int_b^c f = I$  and  $\int_b^c g = J$  then  $f + g$  has integral  $I + J$  on  $[b, c]$ .*

In short  $\int_b^c (f + g) = \int_b^c f + \int_b^c g$  if the right side is defined.

**Proof.** Pick  $\epsilon > 0$ . There is a gauge  $F$  on  $[b, c]$  such that  $\text{Ball}(I, \epsilon)$  contains the Riemann sum for  $f$  on any tagged division of  $[b, c]$  inside  $F$ ; and there is a gauge  $G$  on  $[b, c]$  such that  $\text{Ball}(J, \epsilon)$  contains the Riemann sum for  $g$  on any tagged division of  $[b, c]$  inside  $G$ .

Define  $H(t) = F(t) \cap G(t)$ . Then  $H$  is a gauge on  $[b, c]$ . If  $x_0, \dots, x_n$  is a tagged division of  $[b, c]$  inside  $H$ , then  $x_0, \dots, x_n$  is also inside both  $F$  and  $G$ , so the Riemann sums for  $f$  and  $g$  on  $x_0, \dots, x_n$  are within  $\epsilon$  of  $I$  and  $J$  respectively; thus the Riemann sum for  $f + g$  on  $x_0, \dots, x_n$  is within  $2\epsilon$  of  $I + J$ .  $\square$

**Theorem 17.3.** *Let  $f$  be a function. Let  $a \leq b \leq c$  be real numbers. If  $\int_a^b f = I$  and  $\int_b^c f = J$  then  $f$  has integral  $I + J$  on  $[a, c]$ .*

In short  $\int_a^c f = \int_a^b f + \int_b^c f$  if the right side is defined.

**Proof.** Pick  $\epsilon > 0$ . There is a gauge  $G$  on  $[a, b]$  such that  $\text{Ball}(I, \epsilon)$  contains the Riemann sum for  $f$  on any tagged division of  $[a, b]$  inside  $G$ ; there is a gauge  $H$  on  $[b, c]$  such that  $\text{Ball}(J, \epsilon)$  contains the Riemann sum for  $f$  on any tagged division of  $[b, c]$  inside  $H$ .

I define a new gauge as follows. For  $t < b$  define  $F(t) = \{x \in G(t) : x < b\}$ . For  $t = b$  define  $F(t) = G(t) \cap H(t)$ . For  $t > b$  define  $F(t) = \{x \in H(t) : x > b\}$ .

Say  $x_0, \dots, x_n$  is a tagged division of  $[a, c]$  inside  $F$ . Then  $b \in [x_{k-1}, x_k] \subset F(t_k)$  for some  $k$ ; by construction of  $F$ ,  $t_k$  must equal  $b$ . Now  $x_0, t_1, x_1, \dots, x_{k-1}, t_k, b$  is a tagged division of  $[a, b]$  inside  $F$ , hence inside  $G$ . Thus the Riemann sum  $(x_1 - x_0)f(t_0) + \dots + (b - x_{k-1})f(t_k)$  is within  $\epsilon$  of  $I$ . Similarly the Riemann sum  $(x_k - b)f(t_k) + \dots + (x_n - x_{n-1})f(t_n)$  is within  $\epsilon$  of  $J$ . Add: the Riemann sum  $(x_1 - x_0)f(t_0) + \dots + (x_k - x_{k-1})f(t_k) + \dots + (x_n - x_{n-1})f(t_n)$  is within  $2\epsilon$  of  $I + J$ .  $\square$

**Theorem 17.4.** *Let  $f$  be a function. Let  $b \leq c$  be real numbers. If  $f$  is nonnegative on  $[b, c]$  and  $\int_b^c f = I$  then  $I$  is nonnegative.*

**Proof.** Pick  $\epsilon > 0$ . Select an appropriate gauge  $G$ . By Theorem 14.4, there is an appropriate tagged division of  $[b, c]$ . The corresponding Riemann sum is nonnegative, so  $I \geq -\epsilon$ .  $\square$

## Part VI. Limits

### 18. Convergence and limits

**Definition 18.1.** *Let  $f$  be a function. Then  $f$  converges to  $L$  at  $c$  if the function*

$$x \mapsto \begin{cases} L & \text{if } x = c \\ f(x) & \text{if } x \neq c \end{cases}$$

*is continuous at  $c$ .*

Equivalent terminology:  $f(x)$  **converges to  $L$  as  $x$  approaches  $c$ .**

By Theorem 2.2, there is at most one number  $L$  such that  $f$  converges to  $L$  at  $c$ . If this number exists, it is called **the limit of  $f$  at  $c$** , or **the limit of  $f(x)$  as  $x$  approaches  $c$** , written  $\lim_{x \rightarrow c} f(x)$ . Note that  $f$  is continuous if and only if  $\lim_{x \rightarrow c} f(x) = f(c)$ .

Example:  $\cos(1/x)$  does not converge to 0 as  $x$  approaches 0.

## 19. Limits of sums, products, and compositions

**Theorem 19.1.** *Let  $f$  and  $g$  be functions. If  $\lim_{x \rightarrow c} f(x) = L$  and  $\lim_{x \rightarrow c} g(x) = M$  then  $f(x) + g(x)$  converges to  $L + M$  as  $x$  approaches  $c$ .*

In short  $\lim_{x \rightarrow c} (f(x) + g(x)) = \lim_{x \rightarrow c} f(x) + \lim_{x \rightarrow c} g(x)$  if the right side is defined.

**Proof.** Replace  $f(c)$  by  $L$  and  $g(c)$  by  $M$  to obtain new functions  $a$  and  $b$ . Then  $a$  and  $b$  are continuous, so  $a + b$  is continuous by Theorem 3.1.  $\square$

**Theorem 19.2.** *Let  $f$  and  $g$  be functions. If  $\lim_{x \rightarrow c} f(x) = L$  and  $\lim_{x \rightarrow c} g(x) = M$  then  $f(x)g(x)$  converges to  $LM$  as  $x$  approaches  $c$ .*

**Proof.** Theorem 3.2.  $\square$

**Theorem 19.3.** *Let  $f$  and  $g$  be functions. If  $\lim_{x \rightarrow c} g(x) = L$ , and  $f$  is continuous at  $L$ , then  $f(g(x))$  converges to  $f(L)$  as  $x$  approaches  $c$ .*

In short  $\lim_{x \rightarrow c} f(g(x)) = f(\lim_{x \rightarrow c} g(x))$  if the right side is defined, provided that  $f$  is continuous.

**Proof.** Theorem 3.3.  $\square$

## 20. L'Hôpital's rule

**Theorem 20.1.** *Let  $f$  and  $g$  be real-valued functions differentiable at  $c$ . If  $f(c) = g(c) = 0$ , and  $g'(c) \neq 0$ , then  $f(x)/g(x)$  converges to  $f'(c)/g'(c)$  as  $x$  approaches  $c$ .*

For example,  $\lim_{x \rightarrow 0} (x/\sin x) = 1/1 = 1$ , since  $\sin' = \cos$  and  $\cos 0 = 1 \neq 0$ .

**Proof.** By assumption  $f(x) = f(c) + (x - c)f_1(x) = (x - c)f_1(x)$  where  $f_1$  is continuous at  $c$ . Similarly  $g(x) = (x - c)g_1(x)$  where  $g_1$  is continuous at  $c$ . By assumption  $g_1(c) = g'(c) \neq 0$ , so the function  $x \mapsto f_1(x)/g_1(x)$  is continuous at  $c$ , with value  $f_1(c)/g_1(c)$ . Finally  $f(x)/g(x) = f_1(x)/g_1(x)$  for  $x \neq c$ .  $\square$

**Theorem 20.2.** *Let  $f$  and  $g$  be differentiable real-valued functions. If  $f(c) = g(c) = 0$ , and  $\lim_{x \rightarrow c} (f'(x)/g'(x)) = L$ , then  $f(x)/g(x)$  converges to  $L$  as  $x$  approaches  $c$ .*

**Proof.** Fix a ball  $E$  around  $L$ . There is a ball  $D$  around  $c$  such that  $f'(x)/g'(x) \in E$  for all  $x \in D$  with  $x \neq c$ . In particular,  $g'(x)$  is nonzero for  $x \in D$ . By Theorem 12.1,  $g(y)$  is nonzero for  $y \in D$ .

I will show that  $f(y)/g(y) \in E$  for all  $y \in D$  with  $y \neq c$ . Thus  $f(y)/g(y)$  converges to  $L$  as  $y$  approaches  $c$ .

Given  $y \in D$ ,  $y \neq c$ , consider the function  $h = (x \mapsto f(x)g(y) - f(y)g(x))$ . Notice that  $h$  is differentiable, with  $h'(x) = f'(x)g(y) - f'(y)g'(x)$ .

Now  $h(c) = f(c)g(y) - f(y)g(c) = 0$ , and  $h(y) = f(y)g(y) - f(y)g(y) = 0$ , so there is some  $x$  between  $c$  and  $y$  with  $h'(x) = 0$  by Theorem 12.1. Thus  $f'(x)g(y) = f'(y)g'(x)$ . Both  $g'(x)$  and  $g(y)$  are nonzero, so  $f(y)/g(y) = f'(x)/g'(x) \in E$ .  $\square$

Theorem 20.2 may be used repeatedly. For example:

$$\lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2} = \lim_{x \rightarrow 0} \frac{\sin x}{2x} = \lim_{x \rightarrow 0} \frac{\cos x}{2} = \frac{1}{2}.$$

## 99. Expository notes

Common practice in calculus books is to define continuity using limits. I define limits using continuity; continuity is a simpler concept.

“An open ball around  $c$ ” is substantially easier to read than “for some  $h > 0$ , the set of  $x$  such that  $|x - c| < h$ .”

I use Carathéodory’s definition of the derivative of  $f$ . The point is to give a name to the function  $x \mapsto (f(x) - f(c))/(x - c)$ . I learned about this from an article by Stephen Kuhn in the *Monthly*. It’s also used in the second edition of Apostol’s text.

My proof of Theorem 6.2 uses the formula  $h_1(x) = f_1(x)g(x) + f(c)g_1(x)$ , which is shorter than the (more obvious) formula  $h_1(x) = f_1(x)g(c) + f(c)g_1(x) + (x - c)f_1(x)g_1(x)$ . I was reminded of this simplification by a letter in the *Monthly* from Günter Pickert.

The Heine-Borel theorem follows immediately from Theorem 14.4. See Botsko’s 1987 *Monthly* article for this approach to all the basic completeness theorems. Thanks to Joe Buhler for the reference.

I follow the Kurzweil-Henstock approach to integration. The resulting integral is *more* general than the Lebesgue integral; it is equivalent to the integrals constructed by Denjoy and Perron. There is no need for any technical conditions in the fundamental theorem of calculus, Theorem 16.1; every derivative is integrable. I learned about this from advertisements by Robert G. Bartle in the *Bulletin* and the *Monthly*.